CLAIMS:

1. A room temperature curable organopolysiloxane composition comprising

(A) 100 parts by weight of an organopolysiloxane of the following general formula (1):

$$HO(SiR^1_2O)_nH \qquad (1)$$

wherein R^1 is a substituted or unsubstituted monovalent hydrocarbon radical of 1 to 10 carbon atoms, and n is an integer of at least 10, or an organopolysiloxane of the following general formula (2):

$$(R^2O)_{3-m}\overset{\overset{R^1{}_m}{|}}{Si}O(SiOR^1_2O)_n\overset{\overset{R^1{}_m}{|}}{Si}(OR^2)_{3-m} \qquad (2)$$

wherein R^1 and n are as defined above, R^2 is a substituted or unsubstituted monovalent hydrocarbon radical of 1 to 6 carbon atoms, and m is independently an integer of 0 or 1, or both,

(B) 0.1 to 30 parts by weight of a silane compound having at least two hydrolyzable radicals each attached to a silicon atom in a molecule, the remaining radicals attached to silicon atoms being selected from the group consisting of methyl, ethyl, propyl, vinyl and phenyl, or a partial hydrolyzate thereof or both, and

(C) 0.1 to 10 parts by weight of an organosilicon compound of the following general formula (3):

$$(R^2O)_p\underset{\underset{R^1{}_{3-p}}{|}}{Si}\text{-}R^3\text{-}NH\text{-}R^4\text{-}NH_2 \qquad (3)$$

wherein R^1 and R^2 are as defined above, R^3 is a divalent hydrocarbon radical of 1 to 10 carbon atoms, R^4 is a divalent aromatic ring-bearing hydrocarbon radical of 7 to 10 carbon atoms, and p is an integer of 1 to 3, at least one of the NH and NH_2 radicals being not directly attached to the aromatic ring in R^4.

2. The composition of claim 1 wherein the hydrolyzable radicals in component (B) are selected from among ketoxime, alkoxy, and isopropenoxy radicals.

3. The composition of claim 1 wherein in formula (3), R^2 is methyl or ethyl, and R^3 is methylene, ethylene or propylene.

4. The composition of claim 1 wherein in formula (3), R^4 is selected from the following structures:

$-CH_2-C_6H_4-$ (4),
$-CH_2-C_6H_4-CH_2-$ (5),
$-CH_2-C_6H_4-CH_2-CH_2-$ (6),
$-CH_2-C_6H_4-CH_2-CH_2-CH_2-$ (7),
$-CH_2-CH_2-C_6H_4-$ (8),
$-CH_2-CH_2-C_6H_4-CH_2-$ (9),
$-CH_2-CH_2-C_6H_4-CH_2-CH_2-$ (10),
$-CH_2-CH_2-CH_2-C_6H_4-$ (11) and
$-CH_2-CH_2-CH_2-C_6H_4-CH_2-$ (12).

5. The composition of claim 1 which further comprises a filler.

6. The composition of claim 5 wherein the filler is silica and/or carbon black.

7. The composition of claim 1 which further comprises a condensation reaction catalyst.